



Cornell University
College of Veterinary Medicine

Section of Genetics
Department of Molecular Medicine
C4-140 Veterinary Medical Center
Cornell University
Ithaca, NY 14853-6401
607 253-3610

HERDA REPORT

S+S PARTNERS
700 LOUISIANA ST. SUITE 3100
HOUSTON, TX 77002

Case:	94862-07
Date Received:	7/21/07
Report Date:	7/23/07

Horse: REYS DUAL BADGER	Reg: 4000032
YOB: 2000 Breed: AQHA Sex: M	Alt. ID:
Sire: DUAL REY	Reg: 3258332
Dam: LIPSHY	Reg: 3168649

HERDA Test Result



Result Codes:

- Hr/Hr Affected – Homozygous for HERDA (two copies of HERDA gene).
- N/Hr Carrier – Heterozygous (one normal gene copy and one HERDA gene copy).
- N/N Normal – Homozygous for the normal gene. Does not possess the disease-causing HERDA gene.

Hereditary Equine Regional Dermal Asthenia (HERDA) is inherited as a recessive trait. This means that breedings between two carrier (N/Hr) horses have a 25% chance of producing an affected foal (Hr/Hr). Affected foals generally appear normal at birth and may not show clinical signs of the disorder (fragile, excessively stretchy skin) until they reach one to two years of age. Foals produced by matings between carrier (N/Hr) and normal (N/N) horses always appear to be normal, but 50% of these are expected to be carriers. We therefore recommend DNA testing all offspring produced by carriers of this trait.